

# Ash Whitefly, *Siphoninus phillyreae* (Haliday) (Insecta: Hemiptera: Aleyrodidae: Aleyrodinae)<sup>1</sup>

Ru Nguyen and Avas B. Hamon<sup>2</sup>

## Introduction

The ash whitefly, *Siphoninus phillyreae* (Haliday), is a pest of numerous ornamental and fruit crops, including citrus. It causes severe damage to pear and apple in Europe. Most ash whiteflies in California were found on pomegranate, ash tree, pear, apple, loquat, and citrus. Heavy infestations cause leaf wilt, early leaf drop, and smaller fruit (Bellows et al. 1990).

Discovered in Florida in 2010, if it becomes established it could become a pest of ornamental plants and possibly other crops.



Figure 1. Several life stages of the ash whitefly, *Siphoninus phillyreae* (Haliday), on fallen Bradford pear leaves.

Credits: Lyle J. Buss, University of Florida

## Synonymy

The ash whitefly, *Siphoninus phillyreae*, was described as *Aleyrodes phillyreae* by Haliday (1835), on *Phillyrea latifolia* collected in Dublin, Ireland. It has several synonyms listed in Mound and Halsey (1978).

## Distribution

Native to Europe, the Mediterranean and northern Africa, the ash whitefly is found in numerous countries, including:

**Africa:** Cameroon, Egypt, Ethiopia, Libya, Morocco, Sudan

**Asia:** India, Iran, Israel, Pakistan, Saudi Arabia, Syria

**Europe:** Austria, Corsica, Czech Republic, Cyprus, England, France, Germany, Hungary, Ireland, Italy, Poland, Romania, Spain, Switzerland, Yugoslavia

**North America:** United States (California, Florida)

**Oceania:** Australia (New South Wales, South Australia, Queensland, and Victoria)

(Mound and Halsey 1978, Bellows et al. 1990, Raymond Gill personal communication, Chris Baptista personal communication, Stocks and Hodges 2010).

1. This document is EENY-147, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date September 2003. Revised October 2011. Reviewed December 2014. Visit the EDIS website at <http://edis.ifas.ufl.edu>. This document is also available on Featured Creatures website at <http://entomology.ifas.ufl.edu/creatures>.

2. Ru Nguyen and Avas B. Hamon (retired), Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

In the United States, ash whitefly was first collected in Los Angeles County, California, in 1988, and then spread to other counties. It was later discovered in Arizona, Nevada, and New Mexico. It appeared in Raleigh, North Carolina, in 1993. It is also reported from Arizona, Georgia, Nevada, New Mexico, South Carolina, and Texas. A successful biological control program using a parasitic wasp reduced infestations to undetectable levels (western states) or possibly eliminated the infestation (North Carolina). In 2010, it was discovered in central (Lake Buena Vista) and north-western (Panama City) Florida (Stocks and Hodges 2010).

## Description and Biology

### Adult

The adult appears much like a typical whitefly with a light dusting of white wax. Depending on temperature, females live from 30 to 60 days, while males live an average of nine days. This rapid development time, without the presence of the parasitoid, initially produced numerous generations per year in California, whereas only two to three generations were reported in Egypt. Plus, the ability of all life stages to overwinter on non-deciduous hosts allows a rapid build-up in population at the start of the season (Stocks and Hodges 2010). Winged females lay eggs on the underside of the leaves. When the nymphs emerge, they rarely move far and feed on the plant sap until pupation (Gillespie 2000).



Figure 2. Adult ash whitefly, *Siphoninus phillyreae* (Haliday)  
Credits: Lyle J. Buss, University of Florida

### Pupa

On first observation, the pupal case appears similar to the white armor of a male snow scale. Closer observation with a hand lens reveals a whitefly pupal case with two longitudinal tufts of white wax. The vasiform orifice is surrounded by dark brown derma (inner, thicker layer of the cuticle), with the anal area appearing as a dark brown spot. Lateral areas

of the pupal case are light beige. Depending on the age, lesser or greater amounts of white wax will be present. The dorsal surface has 40 to 50 long, glassy tubercles similar to the cornicles found on aphids. These tubercular structures produce a droplet of glassy wax, and this gives the pupal case an unusual appearance. The longitudinal white tufts of wax obscure some glassy tubercles (R.J. Gill, personal communication). The pupal case size is 0.8 to 1.0 mm long by 0.55 to 0.7 mm wide and is tan or beige in color (Mound 1966, Stocks and Hodges 2010).



Figure 3. Pupal stage of the ash whitefly, *Siphoninus phillyreae* (Haliday), showing glassy, wax droplets.  
Credits: Lyle J. Buss, University of Florida



Figure 4. Puparia (3 tan-colored individuals) and juveniles (translucent nymphs) of the ash whitefly, *Siphoninus phillyreae* (Haliday), on pomegranate.  
Credits: Lyle J. Buss, University of Florida

## Host Plants

The ash whitefly is a polyphagous species found on several plant families (modified from Bellows et al. 1990), including:

### Bignoniaceae:

- *Catalpa X chilopsis*, catalpa hybrid

### Leguminosae:

- *Afzelia* sp., pod mahogany
- *Cercis occidentalis*, western redbud; *C. siliquastrum*, Judas tree

### Lythraceae:

- *Lagerstroemia indica*, crape myrtle

### Magnoliaceae:

- *Liriodendron tulipifera*, tulip tree
- *Magnolia stellata*, star magnolia

### Oleaceae:

- *Fraxinus excelsior*, European ash; *F. latifolia*, Oregon ash; *F. ornus*, flowering ash; *F. syriaca*, Syrian ash; *F. uhdei*, Shamel ash; *F. uhdei* 'Tomlinson', Tomlinson ash; *F. velutina* 'Modesto', Modesto ash; *F. velutina* var. *glabra*, Arizona ash; *F. velutina* var. *coriacea*, western ash
- *Ligustrum* spp., privets
- *Olea africana* (=O. *chrysophylla*), wild olive; *O. europaea*, common olive
- *Phillyrea latifolia* (=P. *media*), a phillyrea
- *Syringa X hyacinthiflora*, a common lilac; *S. laciniata*, cut-leaf lilac; *S. vulgaris*, common lilac

### Punicaceae:

- *Punica granatum*, pomegranate

### Rhamnaceae:

- *Rhamnus alaternus*, buckthorn
- *Ziziphus spina-christi*, crown of thorns

### Rosaceae:

- *Amelanchier denticulata*, serviceberry
- *Chaenomeles X speciosa*, flowering quince

- *Crataegus mollis*, hawthorn; *C. monogyna*, English hawthorn; *C. laevigata* (=C. *oxyacantha*), hawthorn
- *Cydonia oblonga*, quince
- *Eriobotrya deflexa*, golden loquat
- *Heteromeles arbutifolia*, California Christmas berry
- *Malus domestica*, apple; *M. floribunda*, Japanese flowering crab; *M. fusca*, Oregon crabapple; *M. 'Hopa'*, a crabapple; *M. 'Red Jade'*, a crabapple; *M. X scheideckeri*, Scheidecker crabapple
- *Mespilus* sp., mespilus
- *Prunus armeniaca*, apricot; P. X *blireiana*, blue plum hybrid; *P. persica*, peach; *P. salicina*, Santa Rosa plum; *P. virginiana* var. *melanocarpa*, choke cherry
- *Pyracantha* sp., pyracantha
- *Pyrus calleryana*, ornamental pear; *P. communis* (=P. *sativa*), pear; *P. kawakamii*, flowering pear; *P. pyrifolia*, Japanese sand pear

### Rubiaceae:

- *Cephalanthus occidentalis* var. *californicus*, buttonbush

### Rutaceae:

- *Citrus* sp., tangerine; *C. limon*, lemon; *C. sinensis*, navel orange; *C. sinensis*, Valencia orange
- *Fortunella* sp., kumquat.

## Natural Enemies

Ash whitefly has the potential to become a serious pest in new environments. However, it has several natural enemies that can control its populations to under economic thresholds. These natural enemies include (Stocks and Hodges 2010):

### Predators:

- *Clitostethus arcuatus* (Rossi)
- *Menochilus sexmaculatus* (Fabricius)
- *Scymnus pallidivestis* Mulsant

### Parasites:

- *Coccophagus eleaphilus* Silvestri
- *Encarsia inaron* (Walker), *E. partenopea* Masi, *E. formosa* Gahan, *E. galilea* Rivnay, *E. punicae* Hayat, *E. pseudopartenopea* Viggiani and Mazzone, *E. siphonini* Silvestri

- *Eretmocerus corni* Haldeman, *E. siphonini* Viggiani and Battaglia

The California Department of Food and Agriculture imported *Encarsia* sp. from Israel and Italy, and the coccinellid beetle *Clitostethus arcuatus* for the control of ash whitefly in California (Bellows et al. 1990). *Encarsia inaron* was discovered in the ash whitefly infestation in central Florida, while an unidentified parasitoid was associated with the infestation in northwestern Florida.

## Selected References

Bellows TS, Paine TD, Arakawa KY, Meisenbacher C, Leddy P, Kabashimo J. 1990. Biological control sought for ash whitefly. *California Agriculture* 44: 4-6.

CDEA. (2010). Biological control of ash whitefly. *California Department of Food and Agriculture*. <http://www.cdffa.gov/phpps/ipc/biocontrol/83ashwhitefly.htm> (21 September 2010).

Evans GA, Hamon AB. (5 November 2002). Whitefly taxonomic and ecological website: an online interactive catalog of the whiteflies (Hemiptera: Aleyrodidae) of the world and their parasites and predators: [http://www.fsca-dpi.org/homoptera\\_hemiptera/Whitefly/whitefly\\_catalog.htm](http://www.fsca-dpi.org/homoptera_hemiptera/Whitefly/whitefly_catalog.htm) (13 November 2002).

Gillespie PS. (October 2000). A new whitefly for NSW - The ash whitefly. *NSW Agriculture*. <http://www.agric.nsw.gov.au/Hort/ascu/insects/ashwf.htm> (21 September 2010).

Haliday AH. 1835. *Aleyrodes phillyreae*. *Entomology Magazine* 2: 119-120.

Huffman D, Kidd KA. (January 2010). Ash whitefly in North Carolina. *North Carolina State University Insect Notes*. <http://www.ces.ncsu.edu/depts/ent/notes/O&T/trees/note113/note113.html> (21 September 2010).

Mound LA. 1966. A revision of the British Aleyrodidae (Hemiptera: Homoptera). *Bulletin of the British Museum of Natural History, Entomology* 17: 419-420.

Mound LA, Halsey SH. 1978. Whitefly of the World. A Systematic Catalogue of the Aleyrodidae (Homoptera) with Host Plant and Natural Enemy Data. *British Museum of Natural History and John Wiley and Sons, New York, NY*. pp. 192-193.

Paine T, Bellows T, Hoddle M. (May 2009). Ash whitefly. *Center for Invasive Species Research*. [http://cistr.ucr.edu/ash\\_whitefly.html](http://cistr.ucr.edu/ash_whitefly.html) (21 September 2010).

Pickett CH, Pitcairn MJ. 1999. Classical biological control of ash whitefly: factors contributing to its success in California. *Biocontrol* 44: 143-158.

Stocks I, Hodges G. (September 2010). Ash whitefly, *Siphoninus phillyreae* (Haliday), a new exotic whitefly (Hemiptera: Aleyrodidae) in central Florida, and *Encarsia inaron*, its parasitoid (Hymenoptera: Aphelinidae). *Division of Plant Industry*. <http://www.freshfromflorida.com/content/download/23897/486234/siphoninus-phillyreae.pdf> (May 2014).

Viggiani G, Mazzone P. 1980. *Encarsia pseudopartenopea* n.sp., parassita di *Siphoninus phillyreae* (Haliday) (Hom. Aleyrodidae). *Bollettino del Laboratorio di Entomologia Agraria 'Filippo Silvestri'* 37: 9-12.

Viggiani G, Battaglia D. 1983. Le specie italiane del genere *Eretmocerus* Hald. (Hymenoptera: Aphelinidae). *Bollettino del Laboratorio di Entomologia Agraria 'Filippo Silvestri'* 40: 97-101.